

# Detailed three-dimensional anatomic characterization of the human and canine thyroarytenoid and cricothyroid muscles

Eric J. Hunter<sup>1,2,3</sup>

<sup>1</sup> National Center for Voice and Speech, The University of Utah  
Salt Lake City, UT

<sup>2</sup> Department of Bioengineering, The University of Utah  
Salt Lake City, UT

<sup>3</sup> Division of Otolaryngology – Head and Neck Surgery, University of Utah School of Medicine  
Salt Lake City, UT

## Abstract

Detailed muscle information has become increasingly valuable in order to assist the increased complexity of biomechanical models of the larynx. For example, it has become increasingly important to have more details of laryngeal muscle size, direction, muscle structure, muscle shape (e.g., shape of muscle at origin and insertion), and inter-muscle spatial relations. Presented in this report are data of four male and four female canine larynges; specifically presenting details of the intrinsic abductor and adductor musculature of the canine larynx: the posterior cricoarytenoid, the lateral cricoarytenoid and the interarytenoid muscles. Presented are the three-dimensional representations of four to five muscle bundles of each muscle. From this resource, it is expected that biomechanical models of laryngeal mechanisms can take a needed step into realism in order to support and explore clinical phonosurgical therapies. Quantification of vocal fold geometry is necessary for the development of anatomically realistic and consistently defined experimental/computational models of the glottic and subglottic regions. Such models will facilitate the study of the influence of the subglottis in voice production. Updates to this memo can be downloaded at <http://www.vocalfolds.org>.

**Keywords:** laryngeal muscles, posterior cricoarytenoid, lateral cricoarytenoid, interarytenoid, thyroarytenoid, vocal folds

## 1. Introduction

Laryngeal muscles, with the surrounding cartilages and joints, posture the vocal folds via length change and abduction/adduction. Thus, as a primary mechanism behind posturing, laryngeal muscles are key to overall health (ventilation, swallowing, and effort closure of the airway<sup>1</sup>) and voice (vocal onset, self-sustained oscillation, intensity, and pitch<sup>2-7</sup>).

Knowledge of laryngeal structure (e.g., cartilages and soft tissue) and musculature (e.g., intrinsic laryngeal muscles' orientation, strength, and type) is needed to understand the mechanisms of posturing and phonation. Previous studies of laryngeal muscles have largely been whole muscle descriptors, focusing on quantifying average size (i.e., length and cross-sectional area), overall orientation, and mechanical characteristics (e.g., stress-strain relations and contraction times). For example, Cox et al.<sup>8</sup> described the size, length and direction of human and canine cricothyroid (CT) and thyroarytenoid (TA). Finally, whole muscle mechanical characteristics have been reported for many of the laryngeal muscles<sup>9-19</sup>.

Although studies like these have provided a valuable foundation for understanding laryngeal muscles, a disconnect often exists between the findings of previous whole muscle studies (which

use averages as a complete picture of a particular muscle or group of muscles) and the intricacies of a muscle with non-uniform structure, shape, and function, as well as an entire laryngeal system with inter-muscle mechanical dependencies and relationships. For example, whole muscle studies cannot be used to explain why portions of individual laryngeal muscles have specific posturing functions<sup>20-25</sup>. Neither can they be used to explain how to compensate for some medical pathologies or post-operative conditions, which leave only portions of an individual muscle viable for laryngeal control<sup>26;27</sup>. Further, previous whole muscle studies have not addressed inter-muscle spatial relations, which must be known to adequately understand and model such conditions as laryngeal asymmetry, a common symptom of numerous laryngeal pathologies. Thus, specific laryngeal information is particularly important for laryngeal models (of both phonation and posturing), the goal of which is often to lay the foundation to predict vocal injury. If refinements were made to the basic assumptions and the anatomical information which these models are based, the results of small variations in glottal therapy and phonosurgical interventions such as vocal fold medialization could be accurately and non-invasively simulated. Thus, detailed distributed muscle information, which would enhance the understanding of vocal fold mechanics, is essential.

The goal of this manuscript is to present fibre bundle orientations of human and canine CT (in particular, the pars recta, CTR, and the pars oblique, CTO) and TA. Specifically given are laryngeal muscle bundle origin and insertion points in three dimensions, and corresponding average muscle area, approximated with simple cross-sections

## 2. Data

No new data muscle data were collected for the current report. Rather, existing data from both the CT and TA muscles from six human males and three canines were taken from raw data<sup>28</sup>. The larynges were prepared so that the major cartilages and intrinsic muscles remained in the framework. The cricothyroid muscle was exposed (specifically, the CTR and CTO). The TA muscle was exposed by removing the vocal fold mucosa and vocal ligament so that all of the TA muscle fibers were visible. A pin was inserted through both the CT and the TA joints to establish origin points and to keep the joints from moving. A three-dimensional mechanical positioning system with the vernier markings was used to measure ends of the muscle bundles with an accuracy of 0.1mm. For each specimen, the left CT and right TA were dissected. The three-dimensional origin and insertion positions were recorded for each bundle. As a bundle was removed, it was weighed using an electronic balance with 0.1mg accuracy. From this bundle weight and the length from the origin and insertion points, cross-sectional area was calculated. Cox *et al.*<sup>8</sup> published the length of the TA (averaged from the bundles), as well as the area and the angle (direction) in the coordinate system defined in the paper. Also published were the averaged area, angle, and length of the total CT and its two portions, the CTR and CTO.

The raw data was recovered and presented into electronic spreadsheet tables (available with this Technical Memo) for easy access. The main spreadsheet is 'CoxThesisTables-Final.xls'. This spreadsheet contains all of the tables of data<sup>28</sup>. The first tab (AppA) contains CT and TA three dimensional origin and insertion points for multiple muscle bundles for each of the human and canine specimen. The CTR and CTO are labeled. The second tab (AppB) lists the mass of each muscle bundle mass. The third tab (App C) has the length of each muscle bundle, which can also be calculated from AppA. Tab 4 (App D) is the calculated area of each bundle, assuming the

density of muscle tissue of 0.001043 g/mm<sup>3</sup>. The last tab is an equation based sheet which can recalculate the areas for any given density.

### 3. Using the Data

In addition to the tables, three other spreadsheets are provided along with a Matlab script. Two of the spreadsheets are the same information as the first two tabs of the larger spreadsheet presented before (CoxThisisTables\_appA.xls, CoxThisisTables\_appB.xls) while the third spreadsheet (CoxMuscleInfo.xls) contains information about where the data exists in the first two. For example, row three is the information for the first human larynx, with subsequent columns containing numbers used by the Matlab script to find appropriate data about this larynx. Row four is the data for the second larynx, and so on. Row nine is the first canine larynx with the third canine larynx data in row eleven. The Matlab script (CoxData.m) loads the data and plots it in two figures, one presenting the TA muscle and the other presenting the CT muscle.




Figure 1. 3-D rendering of the origin and insertion points of the muscle bundles.

### 4. Accompanying Files

The files mentioned above were packaged in a compressed zip file called NLDR\_0000005\_v10.zip; which contains the following files

File	Description
CoxThisisTables-Final.xls	Full Data of muscle bundles
CoxData.m	Matlab script to create plots of muscle data
CoxMuscleInfo.xls	Used by CoxData.m to load various muscle data from the following two spreadsheets
CoxThisisTables_appA.xls	Subset of full data used by Matlab script
CoxThisisTables_appB.xls	Subset of full data used by Matlab script

## Literature

- (1) Kraus DH, Ali MK, Ginsberg RJ et al. Vocal cord medialization for unilateral paralysis associated with intrathoracic malignancies. *Journal of Thoracic and Cardiovascular Surgery* 1996;111:334-339.
- (2) Hirano M, Vennard W, Ohala J. Regulation of register, pitch and intensity of voice. An electromyographic investigation of intrinsic laryngeal muscles. *Folia Phoniatr (Basel)* 1970;22:1-20.
- (3) Titze IR. The physics of small-amplitude oscillation of the vocal folds. *J Acoust Soc Am* 1988;83:1536-1552.
- (4) Titze IR, Sundberg J. Vocal intensity in speakers and singers. *J Acoust Soc Am* 1992;91:2936-2946.
- (5) Cooke A, Ludlow CL, Hallett N, Selbie WS. Characteristics of vocal fold adduction related to voice onsets. *J Voice* 1997;11:12-22.
- (6) Murry T, Xu JJ, Woodson GE. Glottal configuration associated with fundamental frequency and vocal register. *J Voice* 1998;12:44-49.
- (7) Honda K. Variability analysis of laryngeal muscle activities. In: Titze IR, Scherer RC, eds. *Vocal fold physiology: biomechanics, acoustics and phonatory control*. Denver, CO: The Denver Center for the Performing Arts; 1983:127-137.
- (8) Cox KA, Alipour F, Titze IR. Geometric structure of the human and canine cricothyroid and thyroarytenoid muscles for biomechanical applications. *Ann Otol Rhinol Laryngol* 1999;108:1151-1158.
- (9) Cooper D, Patridge LD, Alipour-Haghighi F. Muscle energetics, vocal efficiency, and laryngeal biomechanics. In: Titze IR, ed. *Vocal fold physiology: frontiers in basic science*. San Diego, CA: Singular Publishing Group; 1993:37-92.
- (10) Cooper DS, Shindo M, Sinha U, Hast MH, Rice DH. Dynamic properties of the posterior cricoarytenoid muscle. *Ann Otol Rhinol Laryngol* 1994;103:937-944.
- (11) Cooper DS, Pinczower E, Rice DH. Thyroarytenoid intramuscular pressures. *Ann Otol Rhinol Laryngol* 1993;102:167-175.
- (12) Perlman AL, Titze IR, Cooper DS. Elasticity of canine vocal fold tissue. *Journal of Speech and Hearing Research* 1984;27:212-219.
- (13) Alipour F, Titze I. Active and passive characteristics of the canine cricothyroid muscles. *J Voice* 1999;13:1-10.
- (14) Alipour F, Titze IR, Hunter EJ, Tayama N. Active and Passive Properties of Canine Abduction/Adduction Laryngeal Muscles. *J Voice* 2005;19:350-359.
- (15) Alipour-Haghighi F, Titze IR, Perlman AL. Tetanic Contraction in Vocal Fold Muscle. *Journal of Speech and Hearing Research* 1989;32:226-231.
- (16) Alipour-Haghighi F, Titze IR. Viscoelastic modeling of canine vocalis muscle in relaxation. *J Acoust Soc Am* 1985;78:1939-1943.
- (17) Alipour-Haghighi F, Titze IR, Durham P. Twitch response in the canine vocalis muscle. *Journal of Speech and Hearing Research* 1987;30:290-294.
- (18) Alipour-Haghighi F, Perlman AL, Titze IR. Tetanic response of the cricothyroid muscle. *Ann Otol Rhinol Laryngol* 1991;100:626-631.
- (19) Alipour-Haghighi F, Titze IR. Elastic models of vocal fold tissues. *J Acoust Soc Am* 1991;90:1326-1331.
- (20) Han Y, Wang J, Fischman DA, Biller HF, Sanders I. Slow tonic muscle fibers in the thyroarytenoid muscles of human vocal folds; a possible specialization for speech. *Anatomical Record* 1999;256:146-157.
- (21) Brandon CA, Rosen C, Georgelis G, Horton MJ, Mooney MP, Sciote JJ. Staining of human thyroarytenoid muscle with myosin antibodies reveals some unique extrafusil fibers, but no muscle spindles. *J Voice* 2003;17:245-254.
- (22) Brandon CA, Rosen C, Georgelis G, Horton MJ, Mooney MP, Sciote JJ. Muscle fiber type composition and effects of vocal fold immobilization on the two compartments of the human posterior cricoarytenoid: a case study of four patients. *J Voice* 2003;17:63-75.
- (23) Sanders I, Han Y, Wang J, Biller H. Muscle spindles are concentrated in the superior vocalis subcompartment of the human thyroarytenoid muscle. [Review] [45 refs]. *J Voice* 1998;12:7-16.
- (24) Sanders I, Rai S, Han Y, Biller HF. Human vocalis contains distinct superior and inferior subcompartments: possible candidates for the two masses of vocal fold vibration. *Ann Otol Rhinol Laryngol* 1998;107:826-833.
- (25) Sanders I, Rao F, Biller HF. Arytenoid motion evoked by regional electrical stimulation of the canine posterior cricoarytenoid muscle. *Laryngoscope* 1994;104:456-462.
- (26) Zealear DL, Billante CR, Courey MS, Sant'Anna GD, Nettekville JL. Electrically stimulated glottal opening combined with adductor muscle botox blockade restores both ventilation and voice in a patient with bilateral laryngeal paralysis. *Ann Otol Rhinol Laryngol* 2002;111:500-506.
- (27) Peretti G, Nicolai P, Redaelli de Zinis LO et al. Endoscopic CO2 laser excision for T1, T1, and T2 glottic carcinomas: cure rate and prognostic factors. *Otolaryngol Head Neck Surg* 2000;123:124-131.
- (28) Cox KA. *Comparative Analysis of the Human and Canine Cricothyroid and Thyroarytenoid Muscles* [ University of Iowa; 1996.

## Acknowledgements

We appreciate Karin Cox for her permission to use the tables from her thesis. Original data collection was supported by National Institutes of Health (NIDCD) project No DC04347; re-creation of the data tables and original matlab script supported by NIDCD project No. DC006801; document preparation, data packaging, and final version of the matlab analysis script supported by NIDCD project No. DC009616. The NRLD is supported by the National Institutes of Health (DC009616). Thanks to Laura M. Hunter for the technical edits.

## Use Agreement

The scripts, images and text are open to use by the public as a service and part of the National Resource of Laryngeal Data (supported by the National Institute of Deafness and other Communicative Disorders, and hosted by the National Center for Voice and Speech). The scripts, images, model and text enclosed in this memo and accompanying this memo are open to use by the public as a service of the NRLD. However, we ask the reader to respect the time and effort put into this manuscript and research. If the text, images, or included scripts are used, the user agrees to reference to this document, the NRLD, and the source of the original data. We also ask the users to consider contacting the original contributors of the data and give them the right of refusal to (1) participate on papers using the data and (2) have their supporting project acknowledged. The user agrees to freely share with the NLDR any extension software build on the data contained.

## Revisions

- 1.0 Eric Hunter: Main document (March 2012)
- 1.1 Eric Hunter: Minor wording fixes (April 2012)